


# Scientific References Describing Applications of GENOVAC Antibodies



## 2007

Catanese MT et al. (2007): High avidity monoclonal antibodies against human scavenger receptor class B type I efficiently block hepatitis C virus infection in the presence of HDL. *J Virol*, 2007 0: JVI.00193-07

Chan CHF et al. (2007): Colorectal Hyperplasia and Dysplasia Due to Human Carcinoembryonic Antigen (CEA) Family Member Expression in Transgenic Mice. *PLoS ONE*, 2/12: e1353

Catanese MT et al. (2007): High avidity monoclonal antibodies against human scavenger receptor class B type I efficiently block hepatitis C virus infection in the presence of HDL. *J Virol*, 2007 0: JVI.00193-07

Schmitter T et al. (2007): Opa Proteins of Pathogenic Neisseriae Initiate Src Kinase-Dependent Lipid Raft-Mediated Uptake via Distinct Human Carcinoembryonic Antigen-Related Cell Adhesion Molecule Isoforms. *Infection and Immunity*, 8:4116-4126

Schreiner A et al. (2007): Junction protein Shrew-1 influences cell invasion and interacts with invasion-promoting protein CD147. *Mol Biol Cell*, 18/4:1272-81

Viswanathan A et al. (2007): Functional expression of N-formyl peptide receptors in human bone marrow-derived mesenchymal stem cells. *Stem Cells*, 25/5:1263-9

Zeisel M et al. (2007): Scavenger receptor class B type I is a key host factor for hepatitis C virus infection required for an entry step closely linked to CD81. *Hepatology*, 46/6:1722-31

## 2006

Avadhanula V et al. (2006): Respiratory viruses augment the adhesion of bacterial pathogens to respiratory epithelium in a viral species- and cell type-dependent manner. *J Virol*, 80/4: 1629–1636

D'Amour K et al (2006): Production of pancreatic hormone-expressing endocrine cells from human embryonic stem cells. *Nat Biotechnol*, 24/1: 1392-1401

Ellies D et al. (2006): Bone density ligand, sclerostin, directly interacts with LRP5 but not LRP5<sup>G171V</sup> to modulate Wnt activity. *J Bone Miner Res*, 21/11:1738-49

Krause S et al. (2006): Blockade of interleukin-13-mediated cell activation by a novel inhibitory antibody to human IL-13 receptor  $\alpha$ 1. *Mol Immunol*: 33/11: 1799-1807

Nöckel J et al (2006): Characterization of gastric adenocarcinoma cell lines established from CEA424/SV40 T antigen-transgenic mice with or without a human CEA transgene. *BMC Cancer*, 6: 57


Pavoni E et al. (2006): Selection, affinity maturation, and characterization of a human scFv antibody against CEA protein. *BMC Cancer*, 6: 41

Weinberg AD et al. (2006): Anti-OX40 (CD134) Administration to nonhuman primates: immunostimulatory effects and toxicokinetic study. *J Immunother*, 29/6: 575-585

## 2005

Bade B et al. (2005): Detection of soluble human granzyme K in vitro and in vivo. *Eur J Immunol.*, 35/10:2940-2948

# Scientific References Describing Applications of GENOVAC Antibodies




- Bade B et al. (2005): Differential expression of the granzymes A, B, K, M and perforin in human peripheral blood lymphocytes. *Int. Immunol.* 17: 1419 - 1428
- Bratke K et al. (2005): Differential expression of human granzymes A, B, and K in natural killer cells and during CD8+ T cell differentiation in peripheral blood. *Eur J Immunol.*, 35/9: 2608-2616
- D'Amour K et al. (2005): Efficient differentiation of human embryonic stem cells to definitive endoderm. *Nat Biotechnol*, 23: 1534-1541
- van Gisbergen K et al. (2005): Interactions of DC-SIGN with Mac-1 and CEACAM1 regulate contact between dendritic cells and neutrophils. *FEBS letters* 579/27: 6159-6168
- Kalina T et al. (2005): Myeloid antigens in childhood lymphoblastic leukemia: clinical data point to regulation of CD66c distinct from other myeloid antigens. *BMC Cancer* 5: 38
- Klaile E et al (2005): CEACAM1 functionally interacts with filamin A and exerts a dual role in the regulation of cell migration. *JCS*, 118: 5513-5524
- Kunii R et al. (2005): Expression of CD13/aminopeptidase N on human gingival fibroblasts and up-regulation upon stimulation with interleukin-4 and interleukin-13. *J Periodontal Res*, 40/2:138-46
- Muenzner P et al (2005): CEACAM engagement by human pathogens enhances cell adhesion and counteracts bacteria-induced detachment of epithelial cells. *JBC*, 170/5: 825-836
- Roessler M et al. (2005): Identification of Nicotinamide *N*-Methyltransferase as a Novel Serum Tumor Marker for Colorectal Cancer. *Clin Cancer Res*, 11/18: 6550-6558

## 2004

- Bharti S (2004): Novel membrane protein shrew-1 targets to Cadherin-mediated junctions in polarized epithelial cells. *Mol Biol Cell*, 15: 397-406.
- Elagoz A et al. (2004): A truncated form of CKb8-1 is a potent agonist for human formyl peptide-receptor-like 1 receptor. *Br J Pharmacol*, 141/1: 37-46.
- Haynes C and BS Schnierle (2004): Retroviral vector targeting through insertion of epidermal growth factor into receptor binding deficient influenza A hemagglutinin results in fusion defective particles. *J Virol Methods*, 120/2: 189-199
- Kammerer R et al. (2004): The tumour suppressor gene CEACAM1 is completely but reversibly downregulated in renal cell carcinoma. *J Pathol*, 204/3: 258-267.
- Kiessling J et al. (2004): Dual targeting of plastid division protein FtsZ to chloroplasts and the cytoplasm. *EMBO reports* 5 /9: 889-894.
- McCaw S et al. (2004): Engulfment of *Neisseria gonorrhoeae*: Revealing distinct processes of bacterial entry by individual carcinoembryonic antigen-related cellular adhesion molecule family receptors. *Infect Immun*, 72/5: 2742-2752.
- Myrtek D et al. (2004): Expression of interleukin-13 receptor  $\alpha$  1-subunit on peripheral blood eosinophils is regulated by cytokines. *Immunology*, 112: 597-604.
- Nanda A et al. (2004): Identification of a binding partner for the endothelial cell surface proteins TEM7 and TEM7R. *Cancer Res*, 64: 8507-8511.
- Schmitter T et al. (2004): Granulocyte CEACAM3 is a phagocytic receptor of the innate immune system that mediates recognition and elimination of human-specific pathogens. *J Exp Med*, 199/1: 35-46.

# Scientific References Describing Applications of GENOVAC Antibodies



## 2003

de Jonge MI et al. (2003): Functional activity of antibodies against the recombinant OpaJ protein from *Neisseria meningitidis*. *Infection and Immunity*, 71/5:2331-2340.

de Jonge MI et al. (2003): Mapping the binding domains on meningococcal Opa proteins for CEACAM1 and CEA receptors. *Mol Microbiol*, 50/3: 1005.

Fahlgren A et al. (2003): Interferon-gamma tempers the expression of carcinoembryonic antigen family molecules in human colon cells: a possible role in innate mucosal defence. *Scand J Immunol*, 58/6:628-641.

## 2002

Billker O et al. (2002): Distinct mechanisms of internalisation of *Neisseria gonorrhoeae* by members of the CEACAM receptor family involving Pac1- and Cdc42-dependent and – independent pathways. *EMBO J*, 21/4: 560-571.

Boulton IC and SD Gray-Owen (2002): Neisserial binding to CEACAM1 arrests the activation and proliferation of CD4+ T lymphocytes. *Nat Immunol*, 3/3:229-236.

Klippel S et al. (2002): Biochemical characterisation of PRV-1, a novel hematopoietic cell surface receptor, which is overexpressed in polycythemia rubra vera. *Blood*, 100/7: 2441-2448.

## 2001

Richter G. et al. (2001): Tumor necrosis factor -alpha regulates the expression of ICOS ligand on CD34+ progenitor cells during differentiation into antigen presenting cells. *J Biol Chem*, 276/49: 45686-93.

Watt SM et al. (2001): Homophilic adhesion of human CEACAM1 involves N-terminal domain interactions: structural analysis of the binding site. *Blood*, 98/5: 1469-1479.

## 2000

Schoelzel S et al. (2000): Carcinoembryonic antigen family members CEACAM6 and CEACAM7 are differentially expressed in normal tissues and oppositely deregulated in hyperplastic colorectal polyps and early adenomas. *Am J Pathol*, 156/2: 595-605

Temerinac S et al. (2000): Cloning of PRV-1, a novel member of the uPAR receptor superfamily, which is overexpressed in polycythemia rubra vera. *Blood*, 95/8: 2569-2576.